

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show S Numbers](#)[Edit S Numbers](#)[Preferences](#)[Cases](#)**Search Results -**

Terms	Documents
dodma	5

Database:

US Patents Full-Text Database
US Pre-Grant Publication Full-Text Database
JPO Abstracts Database
EPO Abstracts Database
Derwent World Patents Index
IBM Technical Disclosure Bulletins

Search:

L1

[Refine Search](#)[Recall Text](#)[Clear](#)**Search History****DATE:** Tuesday, March 19, 2002 [Printable Copy](#) [Create Case](#)

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=USPT; PLUR=YES; OP=ADJ</i>			
<u>L1</u>	dodma	5	<u>L1</u>

END OF SEARCH HISTORY

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show S Numbers](#)[Edit S Numbers](#)[Preferences](#)[Cases](#)**Search Results -**

Terms	Documents
dodma	0

Database:

US Patents Full-Text Database
US Pre-Grant Publication Full-Text Database
JPO Abstracts Database
EPO Abstracts Database
Derwent World Patents Index
IBM Technical Disclosure Bulletins

Search:

L2

[Refine Search](#)[Recall Text](#)[Clear](#)**Search History**DATE: Tuesday, March 19, 2002 [Printable Copy](#) [Create Case](#)

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ</i>			
<u>L2</u>	dodma	0	<u>L2</u>
<i>DB=USPT; PLUR=YES; OP=ADJ</i>			
<u>L1</u>	dodma	5	<u>L1</u>

END OF SEARCH HISTORY

WEST

Generate Collection

Print

L1: Entry 4 of 5

File: USPT

Nov 2, 1999

DOCUMENT-IDENTIFIER: US 5976567 A

TITLE: Lipid-nucleic acid particles prepared via a hydrophobic lipid-nucleic acid complex intermediate and use for gene transfer

Detailed Description Paragraph Right (238):

This example illustrates the encapsulation of plasmid DNA in a lipid vesicles by the detergent dialysis method using different cationic lipids. The dialysis method is as described previously for DODAC (EXAMPLE 1). The amount of plasmid entrapped with different mol % of the various cationic lipids was determined by DEAE Sepharose chromatography (described in EXAMPLE 2). The entrapment efficiency was similar for all cationic lipids tested with approximately 50 to 60% of plasmid DNA. The cationic lipid concentration required in the formulation for optimal plasmid encapsulation was 6.5 % for DOTMA, DSDAC and DODMA-AN in FIG. 41(a); 8% DODAC and DMRIE in 41(b); DCchol in 41(c).

ILE 'HOME' ENTERED AT 18:33:09 ON 19 MAR 2002)

FILE 'MEDLINE, CANCERLIT, EMBASE, BIOSIS, BIOTECHDS, CAPLUS' ENTERED AT
18:34:00 ON 19 MAR 2002

L1 336 S DODMA OR DIMETHYL AMMONIUM CHLORIDE
L2 2758 S CATIONIC LIPID
L3 6 S L2 AND L1
L4 2 DUP REM L3 (4 DUPLICATES REMOVED)